


STEM-ersion



Name _____

Date _____

w/Rational Functions

Anesthesiologist

Eduardo works at a large hospital where his job is to administer anesthesia for surgery. Given a patient's sensitivities, he has to order a variety of anesthesia in various doses. The concentration of the drug in the bloodstream (in mg/L) is modeled by a function where t is minutes since receiving the drug. When the concentration drops below 0.1 mg/L the patient may become conscious. Additional anesthesia must be administered one half hour prior to reaching this level.

Eduardo must write a note to the anesthesiologist who is taking over after his shift. He likes to provide specific directions about who needs more anesthesia and when.

	Patient A	Patient B	Patient C	Patient D	Patient E
Model	$A(t) = \frac{8t}{0.3t^2 + 4.1}$	$B(t) = \frac{7t}{0.4t^2 + 5.3}$	$C(t) = \frac{5.5t}{0.2t^2 + 2.1}$	$D(t) = \frac{4.9t}{0.1t^2 + 3.2}$	$E(t) = \frac{10.2t}{0.3t^2 + 3.9}$
Time Administered	5:04	5:17	5:32	5:51	6:08

Use this space to make any calculations and show work.


Evidence

Conclusion
or Recommendation

Interpret the Evidence. What does it mean?

Analysis
of the evidence

STEM-ersion



Name _____

Date _____

w/ Rational Functions

Anesthesiologist

Eduardo works at a large hospital where his job is to administer anesthesia for surgery. Given different sensitivities, he has to order a variety of anesthesia in various doses. The concentration of the drug in the patient's blood (in mg/L) is modeled by a function where t is minutes since receiving the drug. When the concentration reaches 0.1 mg/L the patient may become conscious. Additional anesthesia must be administered if the concentration falls below this level.

Eduardo must write a note to the anesthesiologist who is taking over care of the patient. He needs to provide specific directions about who needs more anesthesia and when.

	Patient A	Patient B	Patient C	Patient D	Patient E
Model	$A(t) = \frac{8t}{0.3t^2 + 4.1}$	$B(t) = \frac{7t}{0.4t^2 + 5.3}$	$C(t) = \frac{5.5t}{0.2t^2 + 2.1}$	$D(t) = \frac{4.9t}{0.1t^2 + 3.2}$	$E(t) = \frac{10.2t}{0.3t^2 + 3.9}$
Time Administered	5:04	5:17	5:32	5:51	6:08

Use this space to make any calculations and show work.

	Patient A	Patient B	Patient C	Patient D	Patient E
Model	$A(t) = \frac{8t}{0.3t^2 + 4.1}$	$B(t) = \frac{7t}{0.4t^2 + 5.3}$	$C(t) = \frac{5.5t}{0.2t^2 + 2.1}$	$D(t) = \frac{4.9t}{0.1t^2 + 3.2}$	$E(t) = \frac{10.2t}{0.3t^2 + 3.9}$
Time Administered	5:04	5:17	5:32	5:51	6:08
0.1 mg/L in _____ minutes	266.61 minutes	174.92 minutes	274.96 minutes	489.935 minutes	339.96 minutes
Could wake at	-9:30	-8:12	-10:07	-2:01	-11:48
Due for more at	9:00	7:42	9:37	1:31	11:18

Evidence

Conclusion

or Recommendation

Interpret the Evidence. What does it mean?

Analysis
of the evidence

Answers will vary.

Some students may put them in order

The next anesthesiologist would need to administer anesthesia to Patient B first at 7:42, then Patient A at 9:00, then Patient C at 9:37, then Patient E at 11:18, then Patient D at 1:31 if they each remained in surgery.